Recommended Vaccinations for Children

VACCINE	BIRTH	2 MOS	4 MOS	6 MOS	12 MOS	15 MOS	18 MOS	4-6 YRS	11-12 YRS	16-18 YRS
Hepatitis B IMM33	Hep B #1	Нер В #2		Нер В #3						
Pentacel (Dtap/IPV/Hib) Daptacel (Dtap) Quadracel (Dtap/IPV) Adacel (Tdap)		Pentacel #1	Pentacel #2	Pentacel #3		Pentacel #4		Dtap #5 (Dtap/IPV Quadracel combo)	Tdap (Adacel)	
Polio virus		Pentacel #1 (IPV#1)	Pentacel #2 (IPV #2)	Pentacel #3 (IPV#3)				IPV #4 (Dtap/IPV Quadracel combo)		
Hib		Pentacel #1 (Hib #1)	Pentacel #2 (Hib #2)	Pentacel #3 (Hib#3)		Hib #4				
MMR (ProQuad Combo					MMR #1			MMR #2 (optional Proquad combo)		
Varicella (ProQuad Combo)					Varicella #1			Varicella #2 (optional Proquad combo)		
Prevnar		PCV #1	PCV #2	PCV #3		PCV #4				
Meningoccal 4 (ACYW) MENACTRA									Menactra	Menactra
Meningoccal B BEXERO										Bexsero
Hepatitis A					Нер А #1		Нер А #2			
Rotavirus		Rotateq #1	Rotateq #2	Rotateq #3						
HPV									Gardasil	
COVID-19									Pfizer	Pfizer

Gardasil-before 15 years-2 doses (0, 6-12) after 15 years-3 doses (0,2,6)

Bexsero-2 doses beginning ages 16-23 years (0,1)

Flu for children—From 6 months-8 years must have initial 2 dose series (0,1). Given yearly after initial 2 dose series is completed.

COVID-19—Recommended for all children 12 years and up. A second dose is given three weeks after the initial dose.



21SG0125-1323 ©2021 Kettering Health



Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Disease	Vaccine	Disease spread by	Disease symptoms	Disease complications		
Chickenpox	Varicella vaccine protects against chickenpox.	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs)		
Diphtheria	DTaP* vaccine protects against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death		
Hib	Hib vaccine protects against Haemophilus influenzae type b.	Air, direct contact	May be no symptoms unless bacteria enter the blood	Meningitis (infection of the covering around the brain and spinal cord), intellectual disability, epiglottitis (life-threatening infection that can block the windpipe and lead to serious breathing problems), pneumonia (infection in the lungs), death		
Hepatitis A	HepA vaccine protects against hepatitis A.	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders		
Hepatitis B	HepB vaccine protects against hepatitis B.	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer		
Influenza (Flu)	Flu vaccine protects against influenza.	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs)		
Measles	MMR** vaccine protects against measles.	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death		
Mumps	MMR**vaccine protects against mumps.	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord) , encephalitis (brain swelling), inflammation of testicles or ovaries, deafness		
Pertussis	DTaP* vaccine protects against pertussis (whooping cough).	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death		
Polio	IPV vaccine protects against polio.	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death		
Pneumococcal	PCV13 vaccine protects against pneumococcus.	Air, direct contact	May be no symptoms, pneumonia (infection in the lungs)	Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death		
Rotavirus	RV vaccine protects against rotavirus.	Through the mouth	Diarrhea, fever, vomiting	Severe diarrhea, dehydration		
Rubella	MMR** vaccine protects against rubella.	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscar- riage, stillbirth, premature delivery, birth defects		
Tetanus	DTaP* vaccine protects against tetanus.	Exposure through cuts in skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death		

^{*} DTaP combines protection against diphtheria, tetanus, and pertussis.
** MMR combines protection against measles, mumps, and rubella.

Last updated February 2021 • CS322257-A

This material was developed by the CDC. More information is available for free at cdc.gov



8/16/21 11:12 AM

